

Parameter estimation

■ Effect Model (1):

$$Y_{ij} = \mu + A_i + \epsilon_{ij}, \quad \sum J_i A_i = 0$$

Estimation: $\widehat{\mu + A_i} = y_{i\cdot}$ $\hat{\mu} = y_{..}$ $\hat{A}_i = y_{i\cdot} - y_{..}$

Prediction: $\hat{y}_{ij} = \hat{\mu} + \hat{A}_i = y_{i\cdot}$, Residual: $r_{ij} = y_{ij} - y_{i\cdot}$.

■ Effekt Modell (2):

$$Y_{ij} = \mu + A_i + \epsilon_{ij}, \quad A_1 = 0$$

Estimation: $\hat{\mu} = y_{1\cdot}$ $\hat{A}_i = y_{i\cdot} - y_{1\cdot}$

■ Mean Modell: $Y_{ij} = \mu_i + \epsilon_{ij}$

Estimation: $\hat{\mu}_i = y_{i\cdot}$

ANOVA – Regression

- Analysis of variance models can be written as multiple regression models with indicator variables.
- Parameter estimators y_0, y_1, \dots are Least Squares estimators.
- Analysis of variance models are intuitiv, treatment effects can be easily calculated and are uncorrelated.

Berliner Pfannkuchen





Data

Response: Fat absorption of 24 Berliner [g]

Type of Fat	Fat Absorption							Mean
1	164	172	168	177	156	195	172.0	
2	178	191	197	182	185	177	185.0	
3	175	193	178	171	163	176	176.0	
4	155	166	149	164	170	168	162.0	

balanced design: equal replication

Graphical display



R: anova table

```
> mod2=aov(fat~type,data=berliner)
> summary(mod2)
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
type	3	1636.5	545.5	5.4063	0.0069**
Residuals	20	2018.0	100.9		

```
> coef(mod2)
```

(Intercept)	type2	type3	type4
172	13	4	-10

Design matrix

```
> model.matrix(mod2)
```

	(Intercept)	type2	type3	type4
1	1	0	0	0
.....				
6	1	0	0	0
7	1	1	0	0
.....				
12	1	1	0	0
13	1	0	1	0
.....				
18	1	0	1	0
20	1	0	0	1
.....				
24	1	0	0	1

R: Multiple regression I

```
> mod2.r=lm(fat~type,data=berliner)
```

```
> summary(mod2.r)
```

Call:

```
lm(formula = fat ~ type, data = berliner)
```

Residuals:

Min	1Q	Median	3Q	Max
-1.600e+01	-7.000e+00	-1.685e-14	5.250e+00	2.300e+01

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	172.000	4.101	41.943	<2e-16	***
type2	13.000	5.799	2.242	0.0365	*
type3	4.000	5.799	0.690	0.4983	
type4	-10.000	5.799	-1.724	0.1001	

R: Multiple regression II

Residual standard error: 10.04 on 20 degrees of freedom
Multiple R-squared: 0.4478, Adjusted R-squared: 0.365
F-statistic: 5.406 on 3 and 20 DF, p-value: 0.006876

```
> anova(mod2.r)
Analysis of Variance Table
```

Response: fat


	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
type	3	1636.5	545.5	5.4063	0.006876	**
Residuals	20	2018.0	100.9			

Model checking

Modell: $Y_{ij} = \mu + A_i + \epsilon_{ij}, \quad \epsilon_{ij} \sim N(0, \sigma^2)$ i.i.d.

- Normal plot of residuals $r_{ij} = y_{ij} - \bar{y}_i$. To detect Outliers. Normal distribution not crucial in randomized experiments. Nonparametric test: Kruskal-Wallis
- Equal variances: Plot r_{ij} vs y_i .
 $\sigma_{min}^2 < \frac{1}{9}\sigma_{max}^2$ (balanced designs)
log- $\sqrt{-}$ -transformation, weights
- Independent observations: Plot r_{ij} vs time, order more complex model, analysis

Residual plots



Treatment differences

F test significant \Rightarrow There are treatment effects.
Which? How large are the effects?

Treatment differences $y_{i\cdot} - y_{i'\cdot}$.

Fat type 2 – Fat type 1: $185 - 172 = 13$

Fat type 3 – Fat type 1: $176 - 172 = 4$

Fat type 4 – Fat type 1: $162 - 172 = -10$

Standard error of a treatment difference:

$\sqrt{\sigma^2(1/J + 1/J)} = \sqrt{2\sigma^2/J}$, estimated by $\sqrt{2MS_{res}/J}$.

Example: $\sqrt{2 \cdot 100.9/6} = 5.799$

Are Type 2 and 1 significantly different?

t test for $H_0 : A_2 = A_1$

$$t = \frac{y_{2\cdot} - y_{1\cdot}}{\sqrt{2MS_{res}/J}} = \frac{13}{5.799} = 2.242 > 2.086 = t_{0.975, 20}, p = 0.036$$

Confidence interval for Type 2 - Type 1:

$$13 \pm 2.086 \cdot 5.799 = 13 \pm \underbrace{12.097}_{LSD} = (0.9, 25.1)$$

Efficiency of balanced Designs

20 plots in 2 groups
10 + 10

20 plots in 2 groups
1 + 19

Standard error $y_{1.} - y_{2.}$

$$\hat{\sigma} \underbrace{\sqrt{\frac{1}{10} + \frac{1}{10}}}_{0.45}$$

$$\hat{\sigma} \underbrace{\sqrt{1 + \frac{1}{19}}}_{1.03}$$

No big efficiency loss with moderate (2:1) imbalance.

Multiple pairwise comparisons

Are all pairs of treatments different? Is one treatment different from the others? Are there groups of similar treatments? Problem: α_E increases.

- Bonferroni correction for 6 pairwise comparisons:
Significance level: $\alpha_T = 0.05/6$
Critical value: $t_{1-0.05/2.6,20} = 2.927$
Difference between Type 2 and 1 not significant.
- Tukey method for pairwise comparisons:
critical values for the distribution of $\max |y_{i\cdot} - y_{i'\cdot}|$
- Dunnett's method for multiple comparisons with a control group.

Tukey method

Reject $H_0 : A_2 = A_1$, if

$$|t| > \frac{1}{\sqrt{2}} q_{1-\alpha, I, N-I}$$

with $q_{...}$ the quantile of the Studentized Range distribution.


Example: $|t| > \frac{3.958}{\sqrt{2}} = 2.799$.

Type 2 and 1 do not differ significantly.

Tukey Confidence interval for Type 2 - Type 1:

$$13 \pm 2.799 \cdot 5.799 = 13 \pm \underbrace{16.23}_{HSD} = (-3.2, 29.2)$$

R: plot(TukeyHSD(mod2, "type"))



Contrasts

complex comparison: difference between fat types 1 and 4 vs 2 and 3?

Contrast:

$$C = \sum_{i=1}^I \lambda_i A_i \text{ with } \sum \lambda_i = 0$$

C can be estimated by

$$\begin{aligned}\hat{C} &= \sum \lambda_i \hat{A}_i = \sum \lambda_i (y_{i\cdot} - y_{..}) \\ &= \sum \lambda_i y_{i\cdot} - y_{..} \sum \lambda_i = \sum \lambda_i y_{i\cdot}.\end{aligned}$$

Testing of a contrast

Reject $H_0 : \sum_{i=1}^I \lambda_i A_i = 0$, if

$$|t| = \left| \frac{\hat{C}}{\sqrt{MS_{res} \sum \frac{\lambda_i^2}{J}}} \right| > t_{0.975, N-I}$$

Equivalently,

$$F = t^2 = \frac{\hat{C}^2 / \sum \lambda_i^2 / J_i}{MS_{res}} = \frac{SS_C}{MS_{res}}$$

follows a F distribution with 1 and $N - I$ degrees of freedom. SS_C denotes the sum of squares of the contrast C .

Orthogonal contrasts

There are $I - 1$ linearly independent contrasts.

Two contrasts $C_1 = \sum \lambda_i A_i$ and $C_2 = \sum \lambda'_i A_i$ are called **orthogonal**, if $\sum \lambda_i \lambda'_i = 0$.

For balanced designs:

orthogonal contrasts \rightarrow uncorrelated estimates \rightarrow
t tests nearly independent

Partitioning of Treatment Sum of Squares

$$\left(\frac{\hat{C}}{\sqrt{MS_{res} \sum \frac{\lambda_i^2}{J}}} \right)^2 = \frac{J\hat{C}^2 / \sum \lambda_i^2}{MS_{res}} = \frac{SS_C}{MS_{res}} \sim F_{1, N-I}$$

SS_C = Sum of Squares of the contrast C

If C_1, C_2, \dots, C_{I-1} are orthogonal contrasts, then

$$SS_{treat} = SS_{C_1} + SS_{C_2} + \cdots + SS_{C_{I-1}}$$

Summary: Multiple Comparison

n planned , orthogonal con-
trasts ($n \leq I - 1$)

Bonferroni (-Holm) signi-
ficance level α/n

pairwise comparisons

Tukey method

comparison with a control
group

Dunnett's method

complex nonorthogonal or
complex unplanned com-
parisons

Scheffé: critical value
 $\sqrt{(I - 1)F_{I-1,N-I,95\%}}$